



Sumatra-wide assessment of spatiotemporal niche partitioning among small carnivore species

Marsya Christyanti Sibarani¹ · Wido Rizki Albert² · Anton Ario³ · Tomi Ariyanto⁴ · Ryan Avriandy² · Yoan Dinata⁴ · Taufan Kharis⁵ · Dedi Kiswayadi² · Bobby Nopandry⁶ · Wilson Novarino⁷ · Radinal² · Doddy Saputra² · Agus Setiawan⁸ · Sismanto⁹ · Inda D. Solina⁷ · Rikha A. Surya¹⁰ · Made Vasek Wijaya¹¹ · Wai-Ming Wong^{12,13} · Wulan Pusparini¹⁴

Received: 10 June 2021 / Accepted: 23 September 2022 / Published online: 14 October 2022

© The Author(s) under exclusive licence to Deutsche Gesellschaft für Säugetierkunde 2022, corrected publication 2022-

Abstract

Niche partitioning is a result of interspecific competition between closely-related species to allow co-existence. Multiple species of small carnivores co-occur throughout their ranges in Sumatra, but they are among the lesser studied group of mammal species. This study aimed to collate occurrence records of small carnivores, model their island-wide spatial distribution, and assess their spatio-temporal niche partitioning in Sumatra. We collated camera trap records of small carnivores that were mainly bycatch data from widespread tiger surveys. We used Maxent to predict suitable habitat for nine small carnivore species in response to environmental variables, calculated pairwise spatial niche overlap, and then assessed temporal overlap using Kernel density estimation. In total, we detected 16 of the 21 small carnivore species known to occur in Sumatra. We predicted the suitable habitat of nine species that were found in ≥ 20 locations. Species with the smallest extent of predicted suitable habitat were the Malay civet (*Viverra zibetha*) and short-tailed mongoose (*Herpestes brachyurus*). Of 36 pairwise comparisons, five species pairs had high overlaps and four species pairs had low overlap on spatiotemporal niche. High overlaps did not necessarily indicate high competition pressure because these species have different behaviour to allow coexistence, such as food preference and arboreality. Camera trap surveys are commonly conducted for species-specific studies, yet they also yield abundant records of non-target species. We therefore encouraged collaboration among institutions working in the same region to use bycatch data to fill the knowledge gaps in the ecology of other lesser known species.

Keywords Camera trapping · Interspecific competition · Maxent · Niche segregation · Spatial distribution modelling · Species coexistence

Introduction

Multiple species which share similar resource undergo interspecific competition when resources are limited, including within carnivorous mammal community (Petersen et al. 2019). Coexisting carnivores may alter their activity patterns (e.g., Lucherini et al. 2009; Gerber et al. 2012; Chutipong et al. 2017) or use different habitats (St-Pierre et al. 2006; Jennings and Veron 2011; Ramesh et al. 2017) to gain access to more resources, resulting in niche partitioning among the

coexisting species. However, some studies report that carnivores might also co-occur with substantial spatiotemporal niche overlap, indicating differentiation on other niche dimensions such as behaviour and diet preference (Sunarto et al. 2015; Bu et al. 2016).

More than half of mammal species in the order Carnivora belong to small carnivores; this non-taxon group consists of 153 species in nine families distributed globally on every continent except Antarctica and Australasia (Hunter and Barrett 2011). The ecological features of small carnivores as a group are diverse, comprising ground, aquatic, arboreal, and semi-arboreal species, distributed from marine to freshwater ecosystem and the arid desert to moist tropical forests (Schipper et al. 2008). Sumatra, a part of the Indomalayan realm, is home to 21 small carnivore species, among which one species is endemic to Sumatra, i.e. Sumatran hog badger

Handling editor: Francesco Ferretti.

✉ Marsya Christyanti Sibarani
msibarani@wcs.org

Extended author information available on the last page of the article

(*Arctonyx hoevenii*). Sumatra is one of the seven core areas identified for small carnivore conservation (Schreiber et al. 1989).

According to the IUCN Red List of Threatened Species (iucnredlist.org), two small carnivore species in Sumatra are listed as Endangered, three as Near Threatened, four as Vulnerable, and as 12 Least Concern. Across their range in Sumatra, each species coexists with one or more other species of small carnivores. McCarthy and Fuller (2014) recorded six small carnivore species in Bukit Barisan Selatan National Park (NP), southern Sumatra; Holden (2006) reported 14 species in Kerinci Seblat NP, central Sumatra (Holden 2006); in an oil palm plantation, Jennings et al. (2011) found three small carnivore species.

Regardless of the diversity of small carnivores in Sumatra, little is known about their distribution, ecology, and threats as most of the research and conservation effort in this region is more focused on charismatic megafauna, in particular the Sumatran tiger *Panthera tigris sumatrae* (Linkie et al. 2008; Wibisono et al. 2011). Habitat loss is one of the main causes of population declines for small carnivores because most of them are vulnerable to human-modified landscapes (Schipper et al. 2008). In Sumatra, 35.7% of primary forests were lost between 1990 and 2010, and the remaining primary forest cover was only 30% of the land area in 2010 (22% primary degraded forest and 8% primary intact forest) (Margono et al. 2012). Many small carnivore species, however, can thrive in a logged forest (Mathai et al. 2010) and plantations (Jennings et al. 2015) due to high abundance of rodents and other prey. Wildlife trading potentially threatens small carnivore populations. There are only a few reports on small carnivores trading in Sumatra (Shepherd 2008, 2012) and the impact of the trading on wild population of small carnivores remains unknown. Only eight of the 24 small carnivores in Indonesia (3 non-Sumatran species) are listed as protected species under Indonesian law (Ministerial Regulation No. 106 Year 2018 concerning the Updated Protected Species List), and the law enforcement for trading permits and harvest quotas is weak (Shepherd 2008, 2012). Aquatic and semi-aquatic small carnivores are susceptible to the contamination of water bodies (Schipper et al. 2008; Hussain et al. 2011). Habitat loss and unmeasured threats, such as those from wildlife markets, may lead to ‘silent extinctions’. Therefore, it is important to study the status and distribution of small carnivores as the baseline knowledge for conservation management.

In Sumatra, extensive camera trap surveys have been conducted for a few mammal species, such as tiger (O’Brien et al. 2003; Linkie et al. 2006; Puspardini et al. 2018), Sunda clouded leopard (*Neofelis diardi*) (Haidir et al. 2018) and sun bear (*Helarctos malayanus*) (Wong et al. 2012), yet there was also a considerable number of by-catch records of other species. These bycatch records have been used to

assess ecological features of less known species, such as small carnivores (Holden 2006; Cheyne et al. 2010; Pollock et al. 2015) and cats (Puspardini et al. 2014; Sunarto et al. 2015). However, the extents of these studies in Indonesia were usually at a local scale (site-specific). While there have been a number of large-scale studies on Bornean small carnivores (e.g., Kramer-Schadt et al. 2016; Ross et al. 2017; Hearn et al. 2018), little is known about the distribution and how the coexisting small carnivore species segregate niche across space and time in Sumatra. Considering these gaps, we aimed to (i) collate occurrence records of small carnivores from multiple camera trap projects in Sumatra; (ii) predict the spatial distribution patterns of small carnivores; and (iii) assess spatiotemporal niche partitioning among small carnivores in Sumatra.

Methods

Study region

Sumatra is a part of the Sundaland region, which is regarded as one of the global biodiversity hotspots (Myers et al. 2000). The remarkable feature of the island is the distinctive mountainous Bukit Barisan Range that spans the western half of the island, in contrast with the vast plains in the eastern side (Fig. 1). Elevation ranges from sea level to 3,805 m (the peak of Mount Kerinci). The rainfall regions are divided into two: region A in southern part of Sumatra and region B in northern Sumatra. Region A has one high rainfall peak in January and is strongly influenced by wet northwest monsoon in November–March and the dry southeast monsoon in May–September. Region B has two high rainfall peaks in October–November and March–May which are associated with the movement of the inter-tropical convergence zone (Aldrian et al. 2003). Laumonier (1997) divided the bioclimates into the following five categories based on the annual rainfall: subhumid with an annual rainfall of 1000–1500 mm in 1% of Sumatra, humid (1500–2000 mm) in 4.3%, very humid (2000–2500 mm) in 30, superhumid (2500–3000 mm) in 29%, and hyperhumid (> 3000 mm) in 35.7%.

Species occurrence records

We collated small carnivore records from 13 camera trap projects, comprising nine sites/landscapes, conducted by individual researchers and non-profit organisations in collaboration with governmental institutions or private industries across Sumatra (Fig. 1, Table 1). Most of the surveys targeted tigers, except a dataset in Bukit Barisan Selatan NP which targeted terrestrial vertebrates, WMW in Kerinci-Seblat NP for sun bear, and IDS in Southern Solok for small cats. We discarded




Fig. 1 Camera trap survey locations across Sumatra. Most camera trap locations were in forested area (80.2% dryland forest, 13.7% swamp forest, 2.6% agriculture, 2.8% bush-bareland, 0.4% mangrove forest, 0.3% open area)

camera trap records of small carnivores with uncertain species identification and missing time and location information. Of a total 3809 records in the original spreadsheets, we could collect and verify 90.5% of photographs. Of these, we found that there were three empty photos, one unknown small carnivore, and six non-small carnivores photos which were removed. These photographs were then re-verified by Dr. William Duckworth, a small carnivore expert, to ensure reliable species identification.

Predictive spatial distribution modelling of small carnivore species

We used Maxent to create predictive distribution maps of small carnivore species which were detected in at least 20 different camera trap locations. Maxent takes species presences and a set of environmental variables across a geographical space to predict a species's environmental suitability. The output of Maxent modelling is a raster maps

Table 1 Datasets of small carnivore occurrence records and the details of corresponding camera trap surveys

Site	Main target	Survey period	Mean (SD) distance between nearest camera traps (m)	Number of camera trap locations	Survey effort (trap-nights)	Surveyed elevation range (m)	Data source
Ulu Masen PF	Tiger	2012–2013	1769 (766)	151	6504	210–1771	FFI
Gunung Leuser NP	Tiger	2010	2135 (1147)	69	3356	44–2927	GLNP, WCS, Panthera
Gunung Leuser NP	Tiger	2013	1846 (641)	116	5573	106–2996	GLNP, WCS, Panthera
Batang Gadis NP	Tiger	2005–2006	2181 (1372)	16	539	621–1512	BGNP, CI
Restorasi Ekosistem Riau (RER) Concessions, Kampar Peninsula	Tiger	2015	1692 (702)	138	7431	9–33	FFI, RER ¹
Kerinci-Seblat NP	Tiger	2012–2015	1447 (884)	188	9561	273–2273	KSNP, FFI, Panthera
Kerinci-Seblat NP	Sun bears	2009–2011	1921 (410)	126	8813	145–1968	KSNP, WM Wong ²
Solok Selatan, Kerinci-Seblat Landscape	All animals	2015–2018	652 (458)	38	6748	419–1023	ID Solina, Tidar Kerinci Agung Company ³
Berbak Sembilang NP	Tiger	2015–2016	4684 (3999)	15	1791	6–937	B SNP, ZSL
Dangku WR	Tiger	2016	5387 (2555)	15	710	23–937	South Sumatra NCA, ZSL
Bukit Barisan Selatan NP	Terrestrial vertebrate	2010–2015	1385 (24.3)	60	8561	16–320	BBSNP, WCS ⁴
Bukit Barisan Selatan NP	Tiger	2015	2052 (487)	65	9037	164–936	BBSNP, WCS, Panthera ⁵
Bukit Barisan Selatan NP	Tiger	2018	3766 (479)	61	6826		BBSNP, WCS

Sites were sorted from the northernmost to southernmost of Sumatra

NCA Nature Conservation Agency, NP National Park, NR Nature Reserve, PF Protection Forest, WR Wildlife Reserve. References for camera trap survey methods: ¹Avriandy et al. (2016), ²Wong and Linkie (2013), ³Solina et al. (2018), ⁴TEAM Network (2011), ⁵Pusparini et al. (2018)

All cameras were set up without baits

containing a relative index of predicted environmental suitability, where higher values represent a better predicted condition for the species (Phillips et al. 2006). To reduce the effects of spatial autocorrelation, we only used species records that were separated at least 1 km.

We considered eight habitat covariates in the Maxent modelling: elevation, slope, roughness, tree cover, land cover, distance to human-modified habitat, human population density, and annual precipitation. We obtained elevations from SRTM digital elevation model v4.1 at 90 m resolution (Jarvis et al. 2008) and derived slopes and roughness data from the elevation model. We used global tree canopy cover circa 2010 at a 30-m spatial resolution of Hansen et al. (2013) to represent tree cover percentage. We extracted land cover information using a land cover map of Sumatra (MoEF, 2016). The data consisted of 22 land-use types, for which we then combined similar land uses

and excluded water bodies, resulting in six main land-use types; agriculture, bush/bareland, dryland forest, mangrove forest, savannah, and swamp forest. Although the species occurrence data ranged between 2005 and 2018 and we used tree canopy cover and land cover data from only one year, most of the camera trap locations were in protected areas where tree cover is less likely to degrade or lose compared to non-protected areas. Margono et al. (2012) reported that protected areas only accounted for 9% of total forest cover loss in 1990–2010, compared to 65.8% in production forests. Distance to human-modified habitat—agriculture, settlement, and infrastructures of MoEF (2016) land use map—were measured as Euclidean distance from the periphery of modified habitat polygons to grid cells at non-modified habitat (primary forest and secondary forest). This process resulted in all human-modified land uses having a value of 0 m and the highest value occurring in the middle of a large

patch of forest. We used a gridded map of the estimated total number of people per grid-cell at 100 m resolution (WorldPop 2014) to represent human population density. We used average monthly rainfall data at 1 km resolution from WorldClim 2.1 (Fick and Hijmans 2017), and summed them to obtain the annual rainfall data. All raster data were resampled into a resolution of 1 km, i.e., the coarsest resolution of the original raster data, and vector variables were rasterized into a gridded map at the same resolution. We tested for collinearity among variables and removed variables with Pearson's $|\text{r}| > 0.75$ (Kramer-Schadt et al. 2013). As a result, we excluded the roughness and slope (Supporting Information, Table S1). We conducted all spatial data processing and statistical computing using the R programming language version 3.5.2 (R Core Team 2018).

We modelled species distribution using Maxent version 3.4.1 (Phillips et al. 2017). To reduce model overfit, we only used linear, quadratic, and product features for the predictors (Merow et al. 2013). Using simpler features had been suggested to reduce random noise from general species responses to the environment (Syfert et al. 2013). We used tenfold cross-validation to evaluate model predictions for each species. All other model settings remained default (regularization multiplier = 1; maximum number of background points = 10,000). Although our camera trap stations were distributed across Sumatra, they were clustered in only a few sites, mostly in forested areas (Fig. 1). Therefore, we created a sampling bias file by mapping camera trap locations on a 300-m gridded map and assigning a value of 1 on cells within 1 km radius from the camera trap locations. Other cells were given a value of 0.1, following Kramer-Schadt et al. (2013). The result of this background manipulation was that instead of generating background points evenly across Sumatra, Maxent generated more background points around locations where camera traps were placed, and hence approximating the true sampling effort.

Although from camera trap surveys we could obtain detection/non-detection data, our non-detection might occur due to low detection probability instead of true absences. Therefore, we could not use logistic regression, which requires presence/absence data, to predict species distribution. Occupancy modelling allows estimating detection probability from repeated sampling with population closure assumption (Linkie et al. 2007). However, given the differences in survey periods and survey methods (Table 1), our dataset did not meet the closed population assumption and was therefore not suitable for occupancy analysis.

Estimating spatial distribution overlap

To calculate pairwise spatial overlap, we used the function 'calc.niche.overlap' of the R package 'ENMeval' (Muscarella et al. 2014). The function takes two species predicted

distribution maps as inputs and calculates the Schoener's D overlap index based on the habitat suitability value on each grid cells. This overlap index was regarded as a good index to calculate niche overlap based on predictive spatial distribution maps (Rödder and Engler 2011). The index ranged between 0, indicating no overlap, and 1, indicating complete overlap.

Estimating diel activity pattern overlap

At each location, species detections more than 30 min apart were considered to be independent and were used to assess the diel activity patterns of each species (O'Brien et al. 2003; Ridout and Linkie 2009). To estimate the activity pattern overlap, we followed the method developed by Ridout and Linkie (2009) and Linkie and Ridout (2011). Time of species activity recorded by camera trap was treated as circular data ranging from 0 to 2π radians. The coefficient of overlap of species pairs is calculated non-parametrically using Kernel density estimation (Ridout and Linkie 2009). For species pair with less than 75 records each, we chose Δ_1 as the overlap measure. Otherwise, Δ_4 is calculated. Overlap calculations were conducted using the R package 'overlap' (Meredith and Ridout 2014). We used arbitrary cut-off values of > 0.6 for high overlaps and < 0.4 for low overlaps, and categorised values in between as medium overlaps to ease presenting results.

Results

From a total of 75,450 trap nights recorded at 1,058 camera trap locations, we found 16 small carnivore species representing five families (Table 2). Species that are known to occur in Sumatra, yet not recorded in our study, were Javan mongoose (*Herpestes javanicus*), Eurasian otter (*Lutra lutra*), hairy-nosed otter (*Lutra sumatrana*), smooth-coated otter (*Lutrogale perspicillata*), and Malay weasel (*Mustela nudipes*). Species that were frequently recorded and widely distributed across surveyed sites included yellow-throated marten (*Martes flavigula*), banded linsang (*Prionodon linsang*), banded civet (*Hemigalus derbyanus*), masked palm civet (*Paguma larvata*), and binturong (*Arctictis binturong*). Short-tailed mongoose (*Herpestes brachyurus*) and Malay civet (*Viverra zangalunga*) were also frequently recorded, but most of the locations where the two species were recorded (92% and 94%, respectively) were from one site only, i.e. Kampar Peninsula in Central Sumatra.

Predictive distribution models of small carnivore species in Sumatra

The area under the receiver operating characteristic curve (AUC) values of all Maxent models for the nine species was above 0.7, indicating a good model fit to the training data

Table 2 Number of species detections at each study site

Species names	Common names	Red List status	Number of unique camera trap locations with species detections									
			Ulu Masen (151)	Gunung Leuser (185)	Batang Gadis (16)	Kampar Peninsula (138)	Kerinci Seblat (352)	Dangku (15)	Berbak Sembilang (15)	Bukit Barisan Selatan (186)	Total	
Herpestidae												
<i>Herpestes brachyurus</i> *	Short-tailed Mongoose	NT	-	-	-	54	-	1	3	1	1	59
<i>Herpestes javanicus</i>	Javan Mongoose	LC	-	-	-	-	-	-	-	-	-	-
<i>Herpestes semitorquatus</i>	Collared Mongoose	NT	-	1	-	-	-	-	-	1	1	2
Mephitidae												
<i>Mydaus javanensis</i>	Sunda Stink Badger	LC	-	-	-	-	1	-	-	-	11	12
Mustelidae												
<i>Aonyx cinereus</i>	Asian Small-clawed Otter	VU	-	-	-	-	1	-	2	2	2	5
<i>Arctonyx hoevenii</i> *	Sumatran Hog Badger	LC	14	10	-	-	16	-	-	5	5	45
<i>Lutra lutra</i>	Eurasian Otter	NT	-	-	-	-	-	-	-	-	-	-
<i>Lutra sumatrana</i>	Hairy-nosed Otter	EN	-	-	-	-	-	-	-	-	-	-
<i>Lutrogale perspicillata</i>	Smooth-coated Otter	VU	-	-	-	-	-	-	-	-	-	-
<i>Martes flavigula</i> *	Yellow-throated Marten	LC	15	48	4	38	44	1	2	27	27	179
<i>Mustela lutreolina</i>	Indonesian Mountain Weasel	LC	-	1	-	-	-	-	-	-	-	1
<i>Mustela nudpes</i>	Malay Weasel	LC	-	-	-	-	-	-	-	-	-	-
Prionodontidae												
<i>Prionodon linsang</i> *	Banded Linsang	LC	9	32	3	4	31	-	1	14	14	94
Viverridae												
<i>Arctictis binturong</i> *	Binturong	VU	13	4	1	6	13	-	2	15	15	54
<i>Arctogalidia trivirgata</i>	Small-toothed Palm Civet	LC	-	-	-	4	1	-	-	1	1	6
<i>Cynogale bennettii</i>	Sunda Otter Civet	EN	-	-	-	-	-	-	1	1	1	2
<i>Hemigalus derbyanus</i> *	Banded Civet	NT	21	23	-	49	1	1	5	38	38	138
<i>Paguma larvata</i> *	Masked Palm Civet	LC	35	53	10	-	86	-	-	47	47	231
<i>Paradoxurus hermaphroditus</i> *	Common Palm Civet	LC	1	-	-	10	5	7	6	1	1	30
<i>Viverra zangalunga</i> *	Malay Civet	LC	-	-	-	83	-	5	-	-	-	88
<i>Viverricula indica</i>	Small Indian Civet	LC	1	-	-	-	-	-	-	-	-	1

The numbers in brackets under site names indicate the total number of camera trap locations at corresponding sites

*Species included in spatial distribution modelling and spatiotemporal overlap estimation

Table 3 The area under the receiver operating characteristic curve (AUC) and the contribution of environmental layers to the Maxent models of nine small carnivore species

Species	Training AUC	Percent contribution of environmental layers					
		Land cover	Tree cover	Elevation	Distance to modified habitat	Human population	Annual precipitation
Viverridae							
<i>Paguma larvata</i>	0.934	74.5	8.3	9.9	1.9	0	5.4
<i>Paradoxurus hermaphroditus</i>	0.897	78.8	9.9	5	1.2	3.3	1.8
<i>Arctictis binturong</i>	0.901	42.1	53.5	1.9	0.1	0.1	2.3
<i>Hemigalus derbyanus</i>	0.943	22	59.8	5.2	8.6	0	4.3
<i>Viverra zangalla</i>	0.977	91.1	3.6	1.1	2.3	0.1	1.8
Herpestidae							
<i>Herpestes brachyurus</i>	0.985	87.3	6.3	1.4	3.5	0	1.4
Prionodontidae							
<i>Prionodon linsang</i>	0.926	72.2	18.8	5.1	1.4	0.1	2.4
Mustelidae							
<i>Martes flavigula</i>	0.935	54.3	35	5.5	3.3	0.1	1.9
<i>Arctonyx hoevenii</i>	0.971	61.1	13.1	17.4	1.1	0.1	7.2

Response curves of environmental variables are provided in supporting information, Appendix S4

(Table 3). Malay civet and short-tailed mongoose had the smallest extent of predicted suitable habitat and were limited to peat swamp forests (Fig. 2). Predicted suitable habitat of common palm civet (*Paradoxurus hermaphroditus*) mostly covered lowland areas, but was not in highland forests. In contrast, the Sumatran hog badger (*Arctonyx hoevenii*) seemed to be a highland specialist, with most predicted suitable areas covered highland forests and all occurrence records were above 800 m (Supporting Information, Appendix S3). The extents of predicted suitable habitat for the other five small carnivores were mostly in the forested area of Sumatra which dominantly remained in the western part although this pattern may also reflect the sampling efforts which were mostly in forested area.

Diel activity patterns

Of the nine species assessed for activity pattern, five were nocturnal species (banded linsang, banded civet, masked palm civet, common palm civet, and Malay civet). Diurnal species included short-tailed mongoose, Sumatran hog badger, and yellow-throated marten. There was no preference on time of day shown by binturong (*Arctictis binturong*), indicating cathemeral activity pattern (Fig. 3).

Spatio-temporal niche overlaps

Five species pairs showed high spatio-temporal niche overlaps, while there were four species pairs with low overlaps on both spatial and temporal niche (Fig. 4). However, we

also found species pairs with high overlaps on spatial niche, but low temporal overlaps (e.g. between Malay civet and short-tailed mongoose) or vice versa (e.g. between masked palm civet and Malay civet).

Discussion

Our study provides the first island-wide ecological assessment of small carnivores in Sumatra based on a range of camera trap surveys. Other than revealing patterns of niche overlap and partitioning across time and space among small carnivores in Sumatra, our collation of camera trap photographs provided updates on some small carnivore species ranges, some of which have been published elsewhere: Indonesian mountain weasel (*Mustela lutreolina*) in Gunung Leuser NP, which was the first camera trap record for this species (Pusparini and Sibarani 2014) and the first camera trap documentation of a hypopigmented Asian small-clawed otter (*Aonyx cinereus*) in Bukit Barisan Selatan NP (Allen et al. 2019). Previous records of collared mongoose in Sumatra were only from low elevations ≤ 300 m (Holden and Meijaard 2012), but a survey in Gunung Leuser NP recorded this species at a higher altitude of 666 m in July 2013 (Pusparini and Sibarani 2014) and another survey in Bukit Barisan Selatan NP recorded it at 436 m in August 2015 at 7:04 AM (Fig. 5). Small Indian civet (*Viverricula indica*) is widely distributed in mainland Asia and extends to Java, but its extent of occurrence in Sumatra is still




Fig. 2 Predicted distribution maps of nine small carnivore species across Sumatra

unclear (Choudhury et al. 2015). We found one record of small Indian civet (Fig. 5) in its known range in Sumatra, that is in Ulu Masen, at 288 m in a dryland agricultural area, consisting of four occasions between January and February 2013. We recorded a Sunda otter civet (*Cynogale bennettii*), an Endangered small carnivore species,

in a lowland dryland forest at 86 m altitude in 2011 and in a mangrove forest at 19 m in 2015, further confirming its habitat use in lowland forest (Ross et al. 2015).




Fig. 3 Kernel density distribution of activity patterns of nine small carnivores found at >20 locations. The rugs at the bottom of the density curves show the time of the day at which the species were photo-

graphed. Graphs were sorted based on species' diel activity patterns: *N* nocturnal, *D* diurnal, and *C* catheameral

Habitat specialists

Of the nine species modelled for their suitable habitat, we found three species that were predicted to have restricted habitat use: Malay civet, short-tailed mongoose, and Sumatran hog badger. Malay civet and short-tailed mongoose showed a predicted preference for lowland swamp forests in eastern Sumatra and there were only a few records in dryland forests. Previous studies suggest that they are indeed lowland specialists (Jennings and Veron 2011; Duckworth et al. 2016a, b; Ross et al. 2016; Solina et al. 2018). The predictive distribution models by Jennings and Veron (2011) found that evergreen forest is highly suitable habitat for short-tailed mongoose and Malay civet in Southeast Asia. None of the georeferenced occurrence records of the short-tailed mongoose and Malay civet used by Jennings and Veron (2011) was from swamp forests, whereas 95% of short-tailed mongoose and 92% of Malay civet occurrences in our study were from swamp forests. As a result, their resulting prediction maps were markedly differed from those of ours. These contrasting results suggested that species'

predicted suitable habitat might be different when modelled on different scale and that researchers should be cautious when drawing inferences from incomplete data sets. The species occurrence data from our study came from comparable trapping efforts between lowland (< 150 m) dryland and swamp forests (46% trap-nights in dryland forest vs. 40% in swamp forest; 32% camera trap locations in dryland forest vs. 55% in swamp forest), so it might be possible that the Malay civet and short-tailed mongoose in Sumatra preferred lowland swamp forests over dryland forests.

Another habitat specialist small carnivore is the Sumatran hog badger, which was always recorded at higher elevations (> 800 m) in our study. This finding is consistent with the habitat description by Helgen et al. (2008), who suggested that this species occurs in low montane forests to subalpine meadows.

Coexistence mechanisms for small carnivores

The competitive exclusion principle suggests that two species may coexist if they differ on at least one niche



Fig. 4 Species-pair overlap matrix. The numbers on the upper half in the species-pair cells are spatial overlap indices between corresponding species; the numbers on the lower half are temporal overlap indices. Darker shades of green represent higher overlap; darker shades

of red represent lower overlap. Letters in brackets represent family (V Viverridae, H Herpestidae, P Prionodontidae, Mu Mustelidae, Me Mephitidae)

dimension (Hardin 1960). Our findings showed species pairs that had high overlap on one niche dimension and high segregation on another dimension. Malay civet and short-tailed mongoose are two common species in Kampar. Both were mostly found in lowland swamp forests in Sumatra and therefore had high spatial overlap. However, there is almost complete segregation on the temporal niche between the two: the Malay civet is nocturnal, whereas the short-tailed mongoose is mostly active during the day, as found in previous studies (Colón 2002; Jennings et al. 2010; Jennings and Veron 2011).

Species with high niche overlaps on both spatial and diel activity pattern do not necessarily suffer from competition pressure. They may perform segregation on other niche dimensions not assessed in this study, especially their feeding habit. For example, two sympatric mustelid species, yellow-throated marten and Sumatran hog badger, had high niche overlap in their spatial distribution and their activity pattern. However, the yellow-throated marten, a semi-terrestrial species, is generally known as predator of small vertebrates and occasionally feeding on plant matter (Parr and Duckworth 2007), whereas the Sumatran hog badger is strictly terrestrial and feeds on soil

invertebrates, such as earthworms, beetle larvae, and ants, and occasionally non-invertebrates (Helgen et al. 2008).

Study limitations

This study employed camera trap records that mostly came from large mammal surveys. Camera trap installation was done in a way that optimises capture probability for the target species. Therefore, the data may underestimate or overestimate the occurrence of non-target species. First, some small carnivore species in Sumatra are semi-terrestrial and arboreal species which often or rarely come to the ground (Hunter and Barrett 2011); consequently, these species were underrepresented in this study. Second, we only found few records of semi-aquatic species, such as the Sunda otter civet, because most of camera trap stations were not placed close to water bodies. Three of the five non-recorded species, i.e., Eurasian otter (*Lutra lutra*), hairy-nosed otter (*Lutra sumatrana*), and smooth-coated otter (*Lutrogale perspicillata*), are highly water-associated and are hardly recorded in general camera trapping. Another caveat of this study is that sampling effort was biased toward dryland forests (81% of the camera stations); thus the results may underrepresent



Fig. 5 Camera trap photographs of small carnivore species that were found at few camera-trap locations: **a** a collared mongoose in Bukit Barisan Selatan National Park (© BBTNBS, WCS, Panthera), **b** a small Indian civet in Ulu Masen Protection Forest (© BKSDA Aceh,

FFI), **c** a Sunda otter civet in Bukit Barisan Selatan National Park (© BBTNBS, WCS, TEAM Network), and **d** a small-toothed palm civet in Bukit Barisan Selatan National Park (© BBTNBS, WCS, Panthera)

species that prefer non-forest habitat or habitat generalists that are little affected or positively affected by habitat modification and disturbance, such as Javan mongoose, common palm civet, and small Indian civet. Camera traps were usually set up at a relatively high height, 45–50 cm above ground, which may be too high for small carnivores, resulting in missed detection. The datasets we have collated are the most extensive available data for Sumatran small carnivores, but we note that it is unlikely to detect all species present during a survey.

Conclusion and suggestions for future studies

Small carnivores are among the lesser studied mammals, and there is limited research and conservation funding targeting small carnivores compared to their larger cousin carnivores which are often regarded as umbrella and flagship species.

Research efforts for cat species were known to be biased towards larger animals regardless of their threat status (Brodie 2009). Bycatch records of non-target species are useful to infer the distribution and ecology of less known species although it also comes with shortcomings. We, therefore, encourage other researchers and conservation practitioners to collate occurrence data of the other non-target, yet threatened species in Sumatra, such as non-*Panthera* cats, sun bear, dhole (*Cuon alpinus*), Sunda pangolin (*Manis javanica*), Sumatran ground cuckoo (*Carpococcyx viridis*), crested partridge (*Rollulus rouloul*), black partridge (*Melanoperdix niger*), Malay crestless fireback (*Lophura erythrophthalma*), Malay crested fireback (*Lophura rufa*), bronze-tailed peacock-pheasant (*Polyplectron chalcurom*), and great argus (*Argusianus argus*), as well as data deficient species, such as Sumatran striped rabbit (*Nesolagus netscheri*) and Sumatran mountain muntjac (*Muntiacus montanus*). Species population are declining due to increasing extinction threats; therefore, having a better understanding of their current

occurrence and environmental factors affecting them using the best available data is important for their conservation and habitat management. Finally, collaborative research and responsible data-sharing among institutions which conducts camera traps surveys are also crucial to establish a complete regional-scale comparative study.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s42991-022-00315-6>.

Acknowledgements We thank the Indonesian Ministry of Environment and Forestry for granting the permit to undertake our studies. Multiple funding agencies funded our studies: Tropical Ecology Assessment and Monitoring (TEAM) Network, Safe Tiger Fund, Panthera, IUCN-KfW Integrated Tiger Habitat Conservation Programme, Global Environment Facility, and U.S. Fish & Wildlife Service. We thank Hariyo T. Wibisono and Matt Linkie for initiating this collaboration and reviewing the manuscript; William Duckworth for validating species identification and reviewing the manuscript; anonymous reviewers for providing constructive feedback; Dolly Priatna, Adnun Salampessy, Muhammad Iqbal, and Meyner Nusalawo for their contribution in survey design and data collection; Andjar Rafiastanto, Laura D'Arcy, and Agus R. Lembang for supporting the project; and all field staff for the countless hours spent in the forest deploying the camera traps.

Author contributions MS, WP: conceptualization; All authors: methodology; WRA, AA, TA, RA, YD, TK, DK, BN, WN, Radinal, DS, AS, Sismanto, IDS, RAS, MVW, W-MW: Data curation; MS, WP: Formal analysis and investigation; MS: Writing—original draft preparation; All authors: Writing—review and editing; WP: Supervision.

Funding This research used datasets collected from multiple field studies which were funded by different organisations: Tropical Ecology Assessment and Monitoring (TEAM) Network, Safe Tiger Fund, Panthera, IUCN-KfW Integrated Tiger Habitat Conservation Programme, Global Environment Facility, and U.S. Fish & Wildlife Service.

Data availability All data generated or analysed during this study are included in this published article and its supplementary information files.

Code availability Not applicable.

Declarations

Conflict of interest The authors have no relevant financial or non-financial interests to disclose.

Ethical approval This research did not involve the use or handling of animals.

Consent to participate and for publication All authors have declared the consent to participate in and publish the results of this study and that all persons in charge of the datasets have obtained permission from the relevant authorities to carry out the fieldwork and use the data.

References

- Aldrian E, Gates LD, Widodo FH (2003) Variability of Indonesian rainfall and the influence of ENSO and resolution in ECHAM4 simulations and in the reanalyses. Max-Planck-Institut für Meteorologie, Hamburg
- Allen ML, Sibarani MC, Utoyo L (2019) The first record of a wild hypopigmented oriental small-clawed otter (*Aonyx cinereus*). *Ecotropica* 21:201904
- Avriandy R, Gunaryadi D, Pinondang IMR et al (2016) Camera trap report in Restorasi Ekosistem Riau. Fauna & Flora International Indonesia Program
- Brodie JF (2009) Is research effort allocated efficiently for Conservation? Felidae as a global case study. *Biodivers Conserv* 18:2927–2939. <https://doi.org/10.1007/s10531-009-9617-3>
- Bu H, Wang F, McShea WJ et al (2016) Spatial Co-occurrence and activity patterns of mesocarnivores in the temperate forests of Southwest China. *PLoS ONE* 11:1–15. <https://doi.org/10.1371/journal.pone.0164271>
- Cheyne SM, Husson SJ, Chadwick RJ et al. (2010) Diversity and activity of small carnivores of the Sabangau Peat-swamp Forest, Indonesian Borneo. *Small Carniv Conserv* 43:1–7
- Choudhury A, Duckworth JW, Timmins R et al. (2015) *Viverricula indica*. The IUCN Red List of Threatened Species 2015. <https://doi.org/10.2305/IUCN.UK.2015-4.RLTS.T41710A45220632.en>. Accessed 18 Nov 2021
- Chutipong W, Steinmetz R, Savini T, Gale GA (2017) Assessing resource and predator effects on habitat use of tropical small carnivores. *Mammal Res* 62:21–36. <https://doi.org/10.1007/s13364-016-0283-z>
- Colón CP (2002) Ranging behaviour and activity of the Malay civet (*Viverra zangalunga*) in a logged and an unlogged forest in Danum Valley, East Malaysia. *J Zool* 257:473–485. <https://doi.org/10.1017/S0952836902001073>
- Duckworth JW, Hearn AJ, Ross J et al. (2016a) Predicted distribution of short-tailed mongoose *Herpestes brachyurus* (Mammalia: Carnivora: Herpestidae) on Borneo. *Raffles Bull Zool Suppl* 33:132–141
- Duckworth JW, Mathai J, Ross J, Wilting A (2016b) *Herpestes brachyurus*. The IUCN Red List of Threatened Species 2016b: e.T41610A45206655. Accessed 30 Apr 2020
- Fick SE, Hijmans RJ (2017) WorldClim 2: new 1km spatial resolution climate surfaces for global land areas. *Int J Climatol* 37:4302–4315
- Gerber BD, Karpanty SM, Randrianantenaina J (2012) Activity patterns of carnivores in the rain forests of Madagascar: implications for species coexistence. *J Mammal* 93:667–676. <https://doi.org/10.1644/11-mamm-a-265.1>
- Haidir IA, Macdonald DW, Linkie M (2018) Assessing the spatiotemporal interactions of mesopredators in Sumatra's tropical rainforest. *PLoS ONE* 13:e0202876. <https://doi.org/10.1371/journal.pone.0202876>
- Hansen MC, Potapov PV, Moore R et al. (2013) High-resolution global maps of 21st-century forest cover change. *Science* (80-) 342:850–853. <https://doi.org/10.1126/science.1244693>
- Hardin G (1960) The competitive exclusion principle. *Science* (80-) 131:1292–1297
- Hearn AJ, Cushman SA, Ross J et al (2018) Spatio-temporal ecology of sympatric felids on Borneo. Evidence for resource partitioning? *PLoS ONE* 13:1–25. <https://doi.org/10.1371/journal.pone.0200828>
- Helgen KM, Lim NTL, Helgen LE (2008) The hog-badger is not an edentate: systematics and evolution of the genus *Arctonyx* (Mammalia: Mustelidae). *Zool J Linn Soc* 154:353–385. <https://doi.org/10.1111/j.1096-3642.2008.00416.x>
- Holden J (2006) Small carnivores in Central Sumatra. *Small Carniv Conserv* 34&35:34–38
- Holden J, Meijaard E (2012) An orange-coloured collared mongoose *Herpestes semitorquatus* from Aceh, Sumatra, Indonesia. *Small Carniv Conserv* 47:26–29
- Hunter L, Barrett P (2011) *Carnivores of the world*. Princeton University Press, Princeton, NJ

- Hussain SA, Gupta SK, de Silva PK (2011) Biology and ecology of asian small-clawed otter *Aonyx cinereus* (Illiger, 1815): a review. IUCN Otter Spec Gr Bull 28:63–75
- Jarvis A, Reuter HI, Nelson A, Guevara E (2008) Hole-filled SRTM for the globe Version 4, available from the CGIAR-CSI SRTM 90m Database
- Jennings AP, Naim M, Advento AD et al (2015) Diversity and occupancy of small carnivores within oil palm plantations in central Sumatra, Indonesia. Mammal Res 60:181–188. <https://doi.org/10.1007/s13364-015-0217-1>
- Jennings AP, Veron G (2011) Predicted distributions and ecological niches of 8 civet and mongoose species in Southeast Asia. J Mammal 92:316–327. <https://doi.org/10.1644/10-MAMM-A-155.1>
- Jennings AP, Zubaid A, Veron G (2010) Home ranges, movements and activity of the short-tailed mongoose (*Herpestes brachyurus*) on Peninsular Malaysia. Mammalia 74:43–50. <https://doi.org/10.1515/MAMM.2010.001>
- Kramer-Schadt S, Niedballa J, Pilgrim JD et al. (2013) The importance of correcting for sampling bias in MaxEnt species distribution models. Divers Distrib 19:1366–1379. <https://doi.org/10.1111/ddi.12096>
- Kramer-Schadt S, Reinfelder V, Niedballa J et al. (2016) The Borneo carnivore database and the application of predictive distribution modelling. Raffles Bull Zool 2016:18–41
- Laumonier Y (1997) The vegetation and physiography of Sumatra. Kluwer Academic Publisher, Dordrecht
- Linkie M, Chapron G, Martyr DJ et al. (2006) Assessing the viability of tiger subpopulations in a fragmented landscape. J Appl Ecol 43:576–586. <https://doi.org/10.1111/j.1365-2664.2006.01153.x>
- Linkie M, Dinata Y, Nugroho A, Haidir IA (2007) Estimating occupancy of a data deficient mammalian species living in tropical rainforests: sun bears in the Kerinci Seblat region, Sumatra. Biol Conserv 137:20–27. <https://doi.org/10.1016/j.biocon.2007.01.016>
- Linkie M, Haidir IA, Nugroho A, Dinata Y (2008) Conserving tigers *Panthera tigris* in selectively logged Sumatran forests. Biol Conserv 141:2410–2415. <https://doi.org/10.1016/j.biocon.2008.07.002>
- Linkie M, Ridout MS (2011) Assessing tiger-prey interactions in Sumatran rainforests. J Zool 284:224–229. <https://doi.org/10.1111/j.1469-7998.2011.00801.x>
- Lucherini M, Reppucci JJ, Walker RS et al. (2009) Activity pattern segregation of carnivores in the High Andes. J Mammal 90:1404–1409. <https://doi.org/10.1644/09-MAMM-A-002R.1>
- Margono BA, Turubanova S, Zhuravleva I et al. (2012) Mapping and monitoring deforestation and forest degradation in Sumatra (Indonesia) using Landsat time series data sets from 1990 to 2010. Environ Res Lett 7:034010. <https://doi.org/10.1088/1748-9326/7/3/034010>
- Mathai J, Hon J, Juat N et al. (2010) Small carnivores in a logging concession in the Upper Baram, Sarawak, Borneo. Small Carniv Conserv 42:1–9
- McCarthy JL, Fuller TK (2014) Records of small carnivores from Bukit Barisan Selatan National Park, southern Sumatra, Indonesia. Small Carniv Conserv 51:59–63
- Meredith M, Ridout M (2014) Overview of the overlap package
- Merow C, Smith MJ, Silander JA (2013) A practical guide to MaxEnt for modeling species' distributions: what it does, and why inputs and settings matter. Ecography (cop) 36:1058–1069. <https://doi.org/10.1111/j.1600-0587.2013.07872.x>
- Ministry of Environment and Forestry (2016) Spatial Layer: Indonesia Land Use Land Cover 2014
- Muscarella R, Galante PJ, Soley-Guardia M et al. (2014) ENMeval: an R package for conducting spatially independent evaluations and estimating optimal model complexity for ecological niche models. Methods Ecol Evol 5:1198–1205
- Myers N, Mittermeier RA, Mittermeier CG et al. (2000) Biodiversity hotspots for conservation priorities. Nature 403:853–858. <https://doi.org/10.1038/35002501>
- O'Brien TG, Kinnaird MF, Wibisono HT (2003) Crouching tigers, hidden prey: Sumatran tiger and prey populations in a tropical forest landscape. Anim Conserv 6:131–139. <https://doi.org/10.1017/S1367943003003172>
- Parr JWK, Duckworth JW (2007) Notes on diet, habituation and sociality of Yellow-throated Marten *Martes flavigula*. Small Carniv Conserv 36:27–29
- Petersen WJ, Savini T, Steinmetz R, Ngoprasert D (2019) Periodic resource scarcity and potential for interspecific competition influences distribution of small carnivores in a seasonally dry tropical forest fragment. Mamm Biol 95:112–122. <https://doi.org/10.1016/j.mambio.2018.11.001>
- Phillips SJ, Anderson RP, Schapire RE (2006) Maximum entropy modeling of species geographic distributions. Ecol Model 190:231–259. <https://doi.org/10.1016/j.ecolmodel.2005.03.026>
- Phillips SJ, Dudík M, Schapire RE (2017) Maxent software for modeling species niches and distributions (Version 3.4.1). http://biodiversityinformatics.amnh.org/open_source/maxent/. Accessed 11 Jan 2020
- Pollock LJ, Rosauer DF, Thornhill AH et al. (2015) Phylogenetic diversity meets conservation policy: small areas are key to preserving eucalypt lineages. Philos Trans R Soc B Biol Sci. <https://doi.org/10.1098/rstb.2014.0007>
- Pusparini W, Batubara T, Surahmat F et al. (2018) A pathway to recovery: the Critically Endangered Sumatran tiger *Panthera tigris sumatrae* in an “in danger” UNESCO World Heritage Site. Oryx 52:25–34. <https://doi.org/10.1017/S0030605317001144>
- Pusparini W, Sibarani MC (2014) The first record of Indonesian mountain weasel *Mustela lutreolina* from northern Sumatra, Indonesia. Small Carniv Conserv 51:92–95
- Pusparini W, Wibisono HT, Reddy GV, Bharata P (2014) Small and medium sized cats in Gunung Leuser National Park, Sumatra, Indonesia. Cat News 8:4–9
- R Core Team (2018) R: A language and environment for statistical computing. <https://www.r-project.org/>
- Ramesh T, Kalle R, Downs CT (2017) Staying safe from top predators: patterns of co-occurrence and inter-predator interactions. Behav Ecol Sociobiol 71:41
- Ridout MS, Linkie M (2009) Estimating overlap of daily activity patterns from camera trap data. J Agric Biol Environ Stat 14:322–337. <https://doi.org/10.1198/jabes.2009.08038>
- Rödger D, Engler JO (2011) Quantitative metrics of overlaps in Grinnellian niches: advances and possible drawbacks. Glob Ecol Biogeogr 20:915–927. <https://doi.org/10.1111/j.1466-8238.2011.00659.x>
- Ross J, Hearn AJ, Macdonald DW (2017) The Bornean carnivore community: lessons from a little-known guild. In: Macdonald DW, Newman C, Harrington LA (eds) Biology and conservation of musteloids. Oxford University Press
- Ross J, Hearn AJ, Macdonald DW et al. (2016) Predicted distribution of the Malay civet *Viverra zibellina* (Mammalia: Carnivora: Viverridae) on Borneo. Raffles Bull Zool Suppl 33:78–83
- Ross J, Wilting A, Ngoprasert D et al. (2015) *Cynogale bennettii*. In: IUCN Red List Threat. Species. <https://doi.org/10.2305/IUCN.UK.2015-4.RLTS.T6082A45197343.en>. Accessed 22 Nov 2021
- Schipper J, Hoffmann M, Duckworth JW, Conroy J (2008) The 2008 IUCN red listings of the world's small carnivores. Small Carniv Conserv 39:29–34
- Schreiber A, Wirth R, Riffel M, Van Rompaey H (1989) Weasels, civets, mongooses, and their relatives: an action plan for the conservation of mustelids and viverrids. IUCN/SSC Mustelid and Viverrid Specialist Group

- Shepherd CR (2012) Observations of small carnivores in Jakarta wildlife markets, Indonesia, with notes on trade in Javan ferret badger *Melogale orientalis* and on the increasing demand for common palm civet *Paradoxurus hermaphroditus* for civet coffee production. *Small Carniv Conserv* 47:38–41
- Shepherd CR (2008) Civets in trade in Medan, North Sumatra, Indonesia (1997–2001) with notes on legal protection. *Small Carniv Conserv* 38:34–36
- Solina ID, Novarino W, Rizaldi GAJ (2018) Activity pattern and habitat profile of small carnivores in an oil palm landscape. *J Indones Nat Hist* 6:18–27
- St-Pierre C, Ouellet JP, Crête M (2006) Do competitive intraguild interactions affect space and habitat use by small carnivores in a forested landscape? *Ecography (cop)* 29:487–496. <https://doi.org/10.1111/j.0906-7590.2006.04395.x>
- Sunarto S, Kelly MJ, Parakkasi K, Hutajulu MB (2015) Cat coexistence in central Sumatra: ecological characteristics, spatial and temporal overlap, and implications for management. *J Zool* 296:104–115. <https://doi.org/10.1111/jzo.12218>
- Syfert MM, Smith MJ, Coomes DA (2013) The effects of sampling bias and model complexity on the predictive performance of MaxEnt species distribution models. *PLoS ONE* 8:e55158. <https://doi.org/10.1371/journal.pone.0055158>
- TEAM Network (2011) Terrestrial vertebrate monitoring protocol implementation manual, v. 3.1
- Wibisono HT, Linkie M, Guillera-Arroita G et al. (2011) Population status of a cryptic top predator: an island-wide assessment of tigers in Sumatran rainforests. *PLoS ONE*. <https://doi.org/10.1371/journal.pone.0025931>
- Wong W, Leader-Williams N, Linkie M (2012) Quantifying changes in sun bear distribution and their forest habitat in Sumatra. *Anim Conserv* 16:216–223. <https://doi.org/10.1111/j.1469-1795.2012.00587.x>
- WorldPop (2014) Indonesia 100m Population. Alpha version 2010, 2015 and 2020 estimates of numbers of people per pixel (ppp) with national totals adjusted to match UN population division estimates (<http://esa.un.org/wpp/>)
- Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.
- Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

Authors and Affiliations

Marsya Christyanti Sibarani¹ · Wido Rizki Albert² · Anton Ario³ · Tomi Ariyanto⁴ · Ryan Avriandy² · Yoan Dinata⁴ · Taufan Kharis⁵ · Dedi Kiswayadi² · Bobby Nopandry⁶ · Wilson Novarino⁷ · Radinal² · Doddy Saputra² · Agus Setiawan⁸ · Sismanto⁹ · Inda D. Solina⁷ · Rikha A. Surya¹⁰ · Made Vasek Wijaya¹¹ · Wai-Ming Wong^{12,13} · Wulan Pusparini¹⁴

¹ Wildlife Conservation Society - Indonesia Program, Jl. Malabar 1 No. 11, Bogor, West Java 16128, Indonesia

² Fauna & Flora International - Indonesia Programme, Jl. Margasatwa Raya, Komplek Margasatwa Baru No. 7A, Pondok Labu, Cilandak, Jakarta 12450, Indonesia

³ Yayasan Konservasi Indonesia, Jl. Buncit Raya No.22, Pejaten Barat, Pasar Minggu, Jakarta 12510, Indonesia

⁴ Zoological Society of London - Indonesia Program, Jl. Papandayan No. 18 Bogor, West Java 16128, Indonesia

⁵ South Sumatra Nature Conservation Agency (BKSDA Sumsel), Jl. Kol. H. Burlian Km. 6, No. 79, Palembang, Sumatera Selatan 30961, Indonesia

⁶ Batang Gadis National Park, Jl. Willem Iskandar, Sumatera Utara 22915, Indonesia

⁷ Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Andalas, West Sumatra 25163, Indonesia

⁸ Kerinci Seblat National Park, Jl. Basuki Rahmat No. 11, Sungai Penuh, Jambi 22300, Indonesia

⁹ Berbak Sembilang National Park, Jl. Yos Sudarso Km. 4, Jambi 36262, Indonesia

¹⁰ Bukit Barisan Selatan National Park, Jl. Ir. H. Juanda No. 19, Kotaagung, Lampung 35384, Indonesia

¹¹ Riau Ecosystem Restoration, Pangkalan Kerinci, Riau 28300, Indonesia

¹² Durrell Institute of Conservation and Ecology, University of Kent, Canterbury CT2 7NZ, UK

¹³ Panthera, 8 West 40th Street, Floor 18, New York, NY 10018, USA

¹⁴ Wildlife Conservation Research Unit (WildCRU) and Interdisciplinary Center for Conservation Science (ICCS), Department of Zoology, University of Oxford, Oxford, UK